



BY APPOINTMENT TO
HER MAJESTY QUEEN ELIZABETH II
HORSE FOOD MANUFACTURERS
DODSON & HORRELL LIMITED, NANTWICH

DODSON & HORRELL

ANIMAL HEALTH, NUTRITION AND WELL-BEING

FEEDING SUCCESS

FEEDING AND MANAGING THOSE PRONE TO LAMINITIS

The horse or pony prone to laminitis needs a special diet and careful management. Simply watching what you feed isn't enough on its own; you are likely to also have to make some lifestyle changes preventing laminitis is a way of life.



LAMINITIS

All you need to know

What is laminitis?

Laminitis is an extremely painful condition effecting the laminae in the hoof. Laminae are finely structured tissues which bind together the inner hoof wall and the pedal bone. In laminitis the blood flow to the laminae is disrupted, meaning that they weaken and possibly die. If left untreated the pedal bone can become completely unsupported and can rotate within the hoof or sink through the sole. Laminitis is serious and can cause permanent damage so you should always seek veterinary advice. Once your horse and pony has had laminitis they will be more vulnerable to repeat episodes in the future.

What causes laminitis?

There have been many studies on laminitis and it is clear that there is not one single cause. It is now thought that most cases of laminitis are associated with one of three underlying disease states:

- The majority of cases of laminitis are associated with an underlying endocrine disease, commonly Equine Metabolic Syndrome (EMS) or Pituitary Pars Intermedia Dysfunction (PPID or Cushings). Both EMS and PPID are associated with insulin resistance and high blood glucose levels. This is thought to effect blood flow to the laminae in the hooves, possibly through Insulin Growth Factor Receptors.
- Laminitis can also be associated with sepsis or endotoxaemia and this includes grain-induced laminitis due to starch-overload and health problems such as retained placenta.
- Placing excessive weight on one limb for an extended period of time can lead to supporting limb laminitis. This is often due to lameness or pain on the other leg.

Weigh & Fat Score Regularly

Keeping your horse at a healthy weight for their breed and height is an essential part of minimising the risk of laminitis. We recommend fat scoring your horse and weighing them, using our scientifically validated weigh-tape, every fortnight. You should aim for a fat score of between 2.5-3 on a scale of 0-5. Pay particular attention to any regional fat deposits such as a cresty neck, fat pads behind the shoulder or fat-filled eye sockets as these could indicate increased risk of EMS.

Limit Grazing:

Grass is a major source of calories and some horses can eat 5% of their bodyweight as grass, gaining 21kg in a week. To control the amount of grass your horse eats consider strip grazing, using a grazing muzzle, increasing the number of horses in the field or even cutting the grass every week. The risk is higher in the spring and autumn, when grass growth is most rapid, and grass is higher in non-structural carbohydrates such as sugars, starch and fructans. This may increase the risk of laminitis for individuals that are insulin resistant.



Feed Low Calorie Forage:

Horses and ponies prone to laminitis still need plenty of forage in their diets. Restricting forage intake by too much may increase the risk of colic, gastric ulcers and oral stereotypies. Soaking hay can reduce the NSC and calorie content, or alternatively mix hay with lower-calorie oat or barley straw (up to 50:50).

Feed a Suitable Concentrate Feed:

As most horses and ponies prone to laminitis tend to hold weight easily it is unlikely that they will need large concentrate meals, however, it is still very

important that they get a balanced diet, providing all the essential vitamins and minerals for health and well-being. Consider feeding either a low calorie, fibre-based feed that is balanced in vitamins and minerals or a low intake, low calorie balancer.

Feed Small Meals:

Laminitis due to starch-overload is very rare as most horses and ponies prone to laminitis are not fed large amounts of high starch concentrate feeds. It can occur occasionally, for example if a horse or pony breaks into the feed room. You should always ensure feed is kept secure and you feed small concentrate meals.

Introduce all Dietary Changes Gradually:

We recommend you make all changes to the diet gradually; this includes the introduction of forage or pasture. Sudden changes may disrupt the fragile balance of bacteria in the horse's hind-gut which may increase the risk of colic and/or laminitis.

Exercise Daily:

Exercise will help to aid weight loss and can help protect against insulin resistance. Horses should get a minimum of half an hour active walking per day.

Don't Over-Rug:

If your horse has gained weight then winter can be a great time to kick start weight loss. Trace-clipping or using a lighter rug will mean that your horse will burn off some of his excess weight keeping warm.

Provide Digestive Support:

Consider the use of probiotic and prebiotic supplements to help maintain the pH and microflora of the hindgut. Studies show that these can benefit fibre digestion and effect systemic blood inflammatory cytokines.




SAFE & SOUND

Fibre-based feed suitable for those prone to laminitis. Provides additional fibre, vitamins, minerals and essential proteins without promoting weight gain.

Key Features

- Low starch and sugar
- Added Biotin for hoof support
- Herbal blend
- QLC antioxidants to support the body's natural ability to fight against free radical damage

[**CLICK HERE for more information about D&H Safe & Sound**](#)



Feeding rate

400g-600g/100kg of bodyweight per day.

Average scoops used per day (500kg horse): 4-6 stubbs scoops.

1 level Stubbs Scoop approximately contains:
0.5kg/1.1lbs.

Catherine Robertson

“ With four good doers, Safe and Sound is our go-to feed. It has kept my laminitic pony looking lean and, in the vet's words, 'shiny as a seal'. They all love it and it's so easy to use.

The ponies recently broke into the feed room and Safe and Sound was the one they went for! ”



FEEDING GUIDES

Don't forget, more feeding advice and guidelines for feeding for health, behaviour and type of horse can be found on our website. However if you have any concerns over your horses nutritional needs, please don't hesitate to contact us.

EMS

What is Equine Metabolic Syndrome?

Cushings

What is Equine Cushing's Disease?

FEED REVIEW

Want to check what you're currently feeding with us? No problem! Our Feed Review is easy to use in just five minutes, totally free, unbiased and is a personalised feeding evaluation and planning tool for your horse or pony.

FORAGE ANALYSIS

Our Forage Analysis service tests your horse's hay, haylage, grass or straw. Paired with our Feed Review and you've got a complete nutritional consultation and an overview of your horses' total diet.



EQUI-BITES

Fully supplemented vitamin and mineral treats which can be used as a 'top up' for leisure horses and ponies, can be fed straight from the hand.

Key Features

- Low in starch, sugar and calories
- Highly palatable with added herbs
- Contain our patented QLC antioxidant complex
- Suitable for horses and ponies prone to laminitis



WEIGH TAPE

The D&H Weigh Tape is an easy and quick way to measure your horse or ponies weight. Used regularly, weigh tapes can be a consistent option to tracking your horse's weight.

For more help, advice or information
please contact our nutritional helpline advisors



call **01270 782 223**

email **helplineenquiries@dodsonandhorrell.com**

message via **Social Media**

or use the D&H **LiveChat** on our website

Find us on:

 @dodsonandhorrell

 @dodsonhorrell

 @dodsonhorrell

#fedondodsonandhorrell

British 
Equestrian
Team Supplier

DODSON & HORRELL

ANIMAL HEALTH, NUTRITION AND WELL-BEING